



# HDAC1 (Acetyl Lys220) rabbit pAb

| Catalog No         | BYab-00904   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Mouse;Rat  |
| Applications       | WB; ELISA  |
| Gene Name          | HDAC1 RPD3L1   |
| Protein Name       | HDAC1 (Acetyl Lys220)  |
| Immunogen          | Synthesized peptide derived from human HDAC1 (Acetyl Lys220)   |
| Specificity        | This antibody detects endogenous levels of Human, Mouse, Rat HDAC1 (Acetyl Lys220)   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG   |
| Purification       | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution           | WB 1:1000-2000 ELISA 1:5000-20000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | Histone deacetylase 1 (HD1;EC 3.5.1.98)  |
| Observed Band      | 55kD   |
| Cell Pathway       | Nucleus .  |
| Tissue Specificity | Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain.  |
| Function           | negative regulation of transcription from RNA polymerase II promoter, chromatin organization, chromatin remodeling,transcription, regulation of transcription, DNA-dependent, regulation of transcription from RNA polymerase II promoter,protein amino acid deacetylation, anti-apoptosis, positive regulation of cell proliferation, negative regulation of biosynthetic process, positive regulation of biosynthetic process, regulation of specific transcription from RNA polymerase II promoter, positive regulation of specific transcription from RNA polymerase II promoter, negative regulation of specific transcription from RNA polymerase II |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



metabolic process, positive regulation of ge

### **Background**

catalytic activity:Hydrolysis of an N(6)-acetyl-lysine residue of a histone to yield a deacetylated histone.,function:Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.,PTM:Phosphorylation on Ser-421 and Ser-423 promotes enzymatic activity and interactions with NuRD and SIN3 complexes.,PTM:Sumoylated on Lys-444 and Lys-476; which promotes enzymatic activity. Desumoylated by SENP1.,similarity:Belongs to the histone deacetylase family. Type 1 subfamily.,subunit:Part of the core histone deacetylase (HDAC) complex composed of HDAC1, HDAC2, RBBP4 and RBBP7. The core complex associates with MTA2, MBD2, MBD3, MTA1L1, CHD3 and CHD4 to form the nucleosome remodeling and histone deacetylation (NuRD) complex, or with SIN3, SAP18 and SAP30 to form the SIN3 HDAC complex. Component of a BHC histone deacetylase complex that contains HDAC1, HDAC2, HMG20B/BRAF35, AOF2/LSD1, RCOR1/CoREST and PHF21A/BHC80. The BHC complex may also contain ZMYM2, ZNF217, ZMYM3, GSE1 and GTF21. Associates with the 9-1-1 complex; interacts with HUS1. Found in a complex with DNMT3A and HDAC7. Interacts with BCOR, BRMS1L, DAXX, DNMT1, EP300, HCFC1, NFE4, PCAF, PHB2, MIER1, KDM4A, MINT, NRIP1, PRDM6, RERE, SETDB1, SUV39H1, TGIF, TGIF2, UHRF1, UHRF2 and ZNF541. Interacts with the non-histone region of H2AFY. Interacts with BANP, CBFA2T3 and KDM5B. Interacts with SAP30L. Interacts with BANP, CBFA2T3 and KDM5B. Interacts with SAP30L. Interacts with E4F1. Interacts with KFL1 (By similarity). Interacts with SV40 large T antigen., tissue specificity: Ubiquitous, with higher levels in heart, pancreas and testis, and lower levels in kidney and brain.,

## matters needing attention

Avoid repeated freezing and thawing!

#### **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### **Products Images**

Nanjing BYabscience technology Co.,Ltd