



# JNK1 mouse mAb

<b>Catalog No</b>	BYab-04485
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IF
<b>Gene Name</b>	mapk8
<b>Protein Name</b>	
<b>Immunogen</b>	Purified recombinant human JNK1 protein fragments expressed in E.coli.
<b>Specificity</b>	This antibody detects endogenous levels of JNK1 and does not cross-react with related proteins.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse
<b>Purification</b>	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	wb dilution 1:1000 icc dilution 1:100. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Al849689 ;c Jun N terminal kinase 1 ;C-JUN kinase 1 ;c-Jun N-terminal kinase 1 ;EC 2.7.11.24 ;JAK 1A ;JAK1A ;JNK 1 ;JNK 46 ;JNK ;JNK-46 ;JNK1A2 ;JNK21B1/2 ;MAP kinase 8 ;MAPK 8 ;MAPK8 ;Mitogen activated protein kinase 8 ;Mitogen-activated protein kinase 8 ;MK08_HUMAN ;p54 gamma ;PRKM 8 ;PRKM8 ;Protein kinase JNK1 ;Protein kinase, mitogen-activated, 8 ;SAPK 1 ;SAPK gamma ;SAPK1 ;Stress activated protein kinase JNK1 ;Stress-activated protein kinase 1 ;Stress-activated protein kinase JNK1 ;Tyrosine protein kinase JAK1 .
<b>Observed Band</b>	46,54kD
<b>Cell Pathway</b>	Cytoplasm . Nucleus . Cell junction, synapse . In the cortical neurons, predominantly cytoplasmic and associated with the Golgi apparatus and endosomal fraction. Increased neuronal activity increases phosphorylated form at synapses (By similarity). Colocalizes with POU5F1 in the nucleus. .
<b>Tissue Specificity</b>	Brain,Epithelium,Fetal brain,Lung,Pooled,Testis,


**Nanjing BYabscience technology Co.,Ltd**




<b>Function</b>	<p>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,domain:The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.,enzyme regulation:Activated by threonine and tyrosine phosphorylation by either of two dual specificity kinases, MAP2K4 and MAP2K7. Inhibited by dual specificity phosphatases, such as DUSP1.,function:JNK1 isoforms display different binding patterns: beta-1 preferentially binds to c-Jun, whereas alpha-1, alpha-2, and beta-2 have a similar low level of binding to both c-Jun or ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms.,function:Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as JUN, JDP</p>
<b>Background</b>	<p>The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation. Several alternatively spl</p>
<b>matters needing attention</b>	<p>Avoid repeated freezing and thawing!</p>
<b>Usage suggestions</b>	<p>This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.</p>




## Products Images



Western blot detection of JNK1 in C6 and 3T3 cell lysates using JNK1 mouse mAb (1:1000 diluted). Predicted band size: 46,54KDa. Observed band size: 46KDa.



Immunofluorescent analysis of 3T3 cells fixed by anhydrous methanol for 2 h at -20°C and using anti-JNK1 mouse mAb (dilution 1:100).



Western blot detection of JNK1 in CHO-K1 cell lysate(B) and CHO-K1 transfected by JNK1-fragment fusion protein(A) cell lysate using JNK1 mouse mAb (1:2000 diluted). Predicted band size:46,54KDa.Observed band size:46,54KDa.