



AMPK γ1 Mouse mAb(5H5)

Catalog No	BYab-04797
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	IHC;WB
Gene Name	PRKAG1
Protein Name	AMPK γ1
Immunogen	Synthesized peptide derived from human AMPK γ1
Specificity	This antibody detects endogenous levels of AMPK γ1 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.37% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p1:50-200 ,WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	5'-AMP-activated protein kinase subunit gamma-1 (AMPK gamma1) (AMPK subunit gamma-1) (AMPKg)
Observed Band	38kD
Cell Pathway	nucleoplasm,cytosol,membrane,nucleotide-activated protein kinase complex,extracellular exosome,
Tissue Specificity	Fetal liver,Muscle,Testis,
Function	function:AMPK is responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. Also regulates cholesterol synthesis via phosphorylation and inactivation of hydroxymethylglutaryl-CoA reductase and hormone-sensitive lipase. This is a regulatory subunit.,similarity:Belongs to the 5'-AMP-activated protein kinase gamma subunit family.,similarity:Contains 4 CBS domains.,subunit:Heterotrimer of an alpha catalytic subunit, a beta and a gamma
	non-catalytic regulatory subunits. Interacts with FNIP1 and FNIP2.,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. This subunit is one of the gamma regulatory subunits of AMPK. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded Human ProstateTissue using AMPK γ1 Mouse Monoclonal antibody diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Human TestisTissue using AMPK γ1 Mouse Monoclonal antibody diluted at 1:200.

Nanjing BYabscience technology Co.,Ltd