



Aurora A Mouse mAb(2B3)

stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an		
Reactivity Human; Mouse;Rat Applications IHC;WB Gene Name Aurora A Protein Name Aurora A Immunogen Synthesized peptide derived from human Aurora A Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole suin microsit. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1.0 Multiple transcript	Catalog No	BYab-04825
Applications IHC:WB Gene Name Aurora A Protein Name Aurora A Immunogen Synthesized peptide derived from human Aurora A Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle pole in militosis. This gene may play a role in tumor development and progression. A processed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1.	Isotype	IgG
Gene Name Aurora A Protein Name Aurora A Immunogen Synthesized peptide derived from human Aurora A Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle pole chrising to inturnor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1. Multiple transcript	Reactivity	Human; Mouse;Rat
Protein Name Immunogen Synthesized peptide derived from human Aurora A Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in milots)s. This gene may play a role in tumor development and progression. A processed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Applications	IHC;WB
Immunogen Synthesized peptide derived from human Aurora A Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene nearly play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 10. Multiple transcript	Gene Name	Aurora A
Specificity This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene as been found on chromosome 1.0. Multiple transcript	Protein Name	Aurora A
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide. Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1. Multiple transcription.	Immunogen	Synthesized peptide derived from human Aurora A
Source Monoclonal, Mouse Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 10. Multiple transcript	Specificity	This antibody detects endogenous levels of Aurora A at Human, Mouse,Rat
Purification The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.67% sodium azide.
using specific immunogen. Dilution IHC-p1:50-200 ,WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 1.0 Multiple transcript	Source	Monoclonal, Mouse
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 10. Multiple transcript	Dilution	IHC-p1:50-200 ,WB 1:1000-2000
Storage Stability -20°C/1 year Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Concentration	1 mg/ml
Synonyms Observed Band 46kD Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Storage Stability	-20°C/1 year
Cell Pathway Tissue Specificity Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Synonyms	
Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Observed Band	46kD
Function Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Cell Pathway	
Background Aurora A(AURKA) Homo sapiens The protein encoded by this gene is a cell cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Tissue Specificity	
cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript	Function	
	Background	cycle-regulated kinase that appears to be involved in microtubule formation and/or stabilization at the spindle pole during chromosome segregation. The encoded protein is found at the centrosome in interphase cells and at the spindle poles in mitosis. This gene may play a role in tumor development and progression. A processed pseudogene of this gene has been found on chromosome 1, and an unprocessed pseudogene has been found on chromosome 10. Multiple transcript

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of 1) Hela Cell, 2) Mouse Liver Tissue, 3)Rat Liver Tissue Lysate using Aurora A Mouse mAb diluted at 1:2,000.

Nanjing BYabscience technology Co.,Ltd