




## A2AP Polyclonal Antibody

Immunogen Synthesized peptide derived from human protein . at AA range: 390-470  Specificity A2AP Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease: Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function: The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity: Belongs to the serpin family., tissue specificity: Expressed by the liver and secreted in plasma.  Background This gene encodes a member of the serpin family of serine protease inhibitors.		
Reactivity         Human;Rat;Mouse;           Applications         WB;ELISA           Gene Name         SERPINF2 AAP PLI           Protein Name         Alpha-2-antiplasmin (Alpha-2-AP) (Alpha-2-plasmin inhibitor) (Alpha-2-Pl) (Serpin F2)           Immunogen         Synthesized peptide derived from human protein . at AA range: 390-470           Specificity         A2AP Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         54kD           Cell Pathway         Secreted.           Tissue Specificity         Expressed by the liver and secreted in plasma.           Function         disease Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) IMM.2628801, APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity Belongs to the serpin family of s	Catalog No	BYab-05308
Applications WB;ELISA  Gene Name SERPINF2 AAP PLI  Protein Name Alpha-2-antiplasmin (Alpha-2-AP) (Alpha-2-plasmin inhibitor) (Alpha-2-Pl) (Serpin F2)  Immunogen Synthesized peptide derived from human protein . at AA range: 390-470  Specificity A2AP Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM.262850], APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function:The major targets of this inhibitor are plasmin and tryosin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family, sissue specificity:Expressed by the liver and secreted in plasma.  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood citoting pathway, Mutations in this gene result in	Isotype	IgG
Gene Name SERPINF2 AAP PLI  Protein Name Alpha-2-antiplasmin (Alpha-2-AP) (Alpha-2-plasmin inhibitor) (Alpha-2-Pl) (Serpin F2)  Immunogen Synthesized peptide derived from human protein . at AA range: 390-470  Specificity A2AP Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function: The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family, tissue specificity: Expressed by the liver and secreted in plasma.  Background This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in recyulating the blood clotting pathway. Mutations in this gene has a major role in recyulating the blood clotting pathway. Mutations in this gene heas a major role in recyulating the blood clotting pathway. Mutations in this gene hear emuli rice and secreted clotting pathway. Mutations in this gene hear emuli rice and secreted clotting pathway. Mutations in this gene hear emuli rice and secreted clotting pathway. Mutations in this gene hear emuli rice and secreted clotting pathway. Mutations in this gene hear emuli rice and secreted clotting pathway. Mutations in this gene hear emuli rice and secreted clotting path	Reactivity	Human;Rat;Mouse;
Protein Name Alpha-2-antiplasmin (Alpha-2-AP) (Alpha-2-plasmin inhibitor) (Alpha-2-PI) (Serpin F2) Immunogen Synthesized peptide derived from human protein . at AA range: 390-470 Specificity A2AP Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD Cell Pathway Secreted. Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function: The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family, tissue specificity:Expressed by the liver and secreted in plasma.  This gene encodes a member of the serpin family, tissue specificity:Expressed by the liver and secreted in plasman, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulation the blood clotting pathway. Mutations in this gene result in requisition the blood clotting pathway. Mutations in this gene result in requisition the blood clotting pathway. Mutations in this gene result in requisition to the proteins. Consequently, the proper function of this gene has a major role in requisition the blood clotting pathway. Mutations in this gene result in the proteins.	Applications	WB;ELISA
F2   Synthesized peptide derived from human protein . at AA range: 390-470	Gene Name	SERPINF2 AAP PLI
Specificity  A2AP Polyclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  54kD  Cell Pathway  Secreted.  Tissue Specificity  Expressed by the liver and secreted in plasma.  function  disease: Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function: The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity: Belongs to the serpin family, tissue specificity: Expressed by the liver and secreted in plasma.  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major rotein is erequilating the blood clotting pathway.	Protein Name	Alpha-2-antiplasmin (Alpha-2-AP) (Alpha-2-plasmin inhibitor) (Alpha-2-PI) (Serpin F2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Immunogen	Synthesized peptide derived from human protein . at AA range: 390-470
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell Pathway         Cell Pathway       Secreted.         Tissue Specificity       Expressed by the liver and secreted in plasma.         Function       disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family. tissue specificity:Expressed by the liver and secreted in plasma.         Background       This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotifup pathway. Mutations in this gene result in	Specificity	A2AP Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  54kD  Cell Pathway  Secreted.  Tissue Specificity  Expressed by the liver and secreted in plasma.  Function  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family, tissue specificity:Expressed by the liver and secreted in plasma.  Background  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis. function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin, similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.  Background This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       -20°C/1 year         Observed Band       54kD         Cell Pathway       Secreted.         Tissue Specificity       Expressed by the liver and secreted in plasma.         Function       disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis., function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity:Belongs to the serpin family., tissue specificity:Expressed by the liver and secreted in plasma.,         Background       This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis, function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  Background This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms  Observed Band 54kD  Cell Pathway Secreted.  Tissue Specificity Expressed by the liver and secreted in plasma.  Function disease: Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis., function: The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity: Belongs to the serpin family., tissue specificity: Expressed by the liver and secreted in plasma.,  Background This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Concentration	1 mg/ml
Synonyms  Observed Band  54kD  Cell Pathway  Secreted.  Tissue Specificity  Expressed by the liver and secreted in plasma.  Function  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Purity	≥90%
Observed Band  Cell Pathway  Secreted.  Tissue Specificity  Expressed by the liver and secreted in plasma.  Function  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  Background  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Storage Stability	-20°C/1 year
Tissue Specificity  Expressed by the liver and secreted in plasma.  Gisease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis., function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin., similarity:Belongs to the serpin family., tissue specificity:Expressed by the liver and secreted in plasma.,  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Synonyms	
Tissue Specificity  Expressed by the liver and secreted in plasma.  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Observed Band	54kD
Function  disease:Defects in SERPINF2 are the cause of alpha-2-plasmin inhibitor deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Cell Pathway	Secreted.
deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed by the liver and secreted in plasma.,  This gene encodes a member of the serpin family of serine protease inhibitors. The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Tissue Specificity	Expressed by the liver and secreted in plasma.
The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in regulating the blood clotting pathway. Mutations in this gene result in	Function	deficiency (APLID) [MIM:262850]. APLID is an autosomal recessive disorder resulting in severe hemorrhagic diathesis.,function:The major targets of this inhibitor are plasmin and trypsin, but it also inactivates chymotrypsin.,similarity:Belongs to the serpin family.,tissue specificity:Expressed
	Background	The protein is a major inhibitor of plasmin, which degrades fibrin and various other proteins. Consequently, the proper function of this gene has a major role in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	hemorrhagic diathesis. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2009],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images