



PX11A Polyclonal Antibody

Catalog No BYab-05905 Isotype IgG Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name PEX11A PEX11 Protein Name Peroxisomal membrane protein 11A (HsPEX11p) (28 kDa peroxisomal integral membrane protein) (PMP28) (Peroxin-11A) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog aipha) (PEX11-alpha) Immunogen Synthesized peptide derived from human protein , at AA range: 110-190 Specificity PX11A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms ZrkD Ceil Pathway Peroxisome membrane ; Multi-pass membrane protein . Tissue Specificity Breast, Liver, Lung, PCR rescued clones, Function function:May be involved in peroxisomal proliferation and may regulate peroxisomes division. M		
ReactivityHuman;Rat:MouseApplicationsWB;ELISAGene NamePEX11A PEX11Protein NamePeroxisomal membrane protein 11A (HsPEX11p) (28 kDa peroxisomal integral membrane protein) (PMP28) (Peroxin-11A) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog alpha) (PEX11-alpha)ImmunogenSynthesized peptide derived from human protein . at AA range: 110-190SpecificityPX11A Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsClasse membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisome division. May mediate binding of coatomer proteins to the peroxisome all interacts with COPB2 and COPA (By stimilarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane proliferation. This gene product in teracts with peroxisome antiperane protein 19 and may respond to outifies thing of coatomer proteins to the peroxisome antiperator of the reproxisome antiperance and proliferation. This gene product interacts with peroxisome maintenance. Alternatively spliced transcript variants encoding multiple isoforms have been <td>Catalog No</td> <td>BYab-05905</td>	Catalog No	BYab-05905
ApplicationsWB;ELISAGene NamePEX11A PEX11Protein NamePeroxisomal membrane protein 11A (HsPEX11p) (28 KDa peroxisomal integral membrane protein) (PMP28) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog alpha) (PEX11-alpha)ImmunogenSynthesized peptide derived from human protein . at AA range: 110-190SpecificityPX11A Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit.IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsEnvisione membrane ; Multi-pass membrane protein .Tissue SpecificityBreast_Liver, Lung, PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisomal membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges to the peroxisoma membrane., PTM: Seems not to be N-glycosylated, similarity. Belonges <td>Isotype</td> <td>lgG</td>	Isotype	lgG
Gene Name PEX11A PEX11 Protein Name Peroxisomal membrane protein 11A (HsPEX11p) (28 kDa peroxisomal integral membrane protein) (PMP28) (Peroxin-11A) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog alpha) (PEX11-alpha) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity PX11A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms End 27kD Cell Pathway Peroxisome membrane ; Multi-pass membrane protein . Function function:May be involved in peroxisomal proliferation and may regulate peroxisomal membranePTM:Seems not to be N-glycosylated, similarity.Belongs to the peroxisomal membranePTM:Seems not to be N-glycosylated, similarity.Belongs to the peroxisomal membranePTM:Seems not to be N-glycosylated, similarity.Belongs to the peroxisomal membranePTM:Seems not to be N-glycosylated, similarity.Belongs to the peroxisomal membranePTM:Seems not to be N-glycosylated, similarity.Belongs to the peroxi	Reactivity	Human;Rat;Mouse
Protein Name Peroxisomal membrane protein 11A (HsPEX11p) (28 kDa peroxisomal integral membrane protein) (PMP28) (Peroxin-11A) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog alpha) (PEX11-alpha) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity PX11A Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Observed Band 27kD Cell Pathway Peroxisome membrane ; Multi-pass membrane protein . Tissue Specificity Breast, Liver, Lung, PCR rescued clones, Function function:May be involved in peroxisomal proliferation and may regulate peroxisome divison. May mediate binding of coatomer proteins to the peroxisome divison. May mediate binding of coatomer proteins to the peroxisome divison. May mediate binding of coatomer proteins to the peroxisome divison. May mediate binding of coatomer proteins to the peroxisome divison. May mediate binding of coatomer proteins to the peroxisome divison. May	Applications	WB;ELISA
ImmunogenSynthesized peptide derived from human protein . at AA range: 110-190SpecificityPX11A Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomea division. May mediate binding of coatomer proteins to the peroxisomea division. May mediate binding of coatomer proteins to the 	Gene Name	PEX11A PEX11
SpecificityPX11A Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit, IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCollObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomal membrane.Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomal membrane.BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome admembrane and the peroxisome admembrane protein 19 and may respond to outside stimuli to increase peroxisome admembrane protein in peroxisome and membrane protein in peroxisome admembrane	Protein Name	Peroxisomal membrane protein 11A (HsPEX11p) (28 kDa peroxisomal integral membrane protein) (PMP28) (Peroxin-11A) (Peroxisomal biogenesis factor 11A) (Protein PEX11 homolog alpha) (PEX11-alpha)
FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCollectionObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in neroxisomal membrane and peroxisome and proliferation. This gene provisione and manter and may regulate provisione and 	Immunogen	Synthesized peptide derived from human protein . at AA range: 110-190
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 27kD Cell Pathway Peroxisome membrane ; Multi-pass membrane protein . Tissue Specificity Breast, Liver, Lung, PCR rescued clones, Function function:May be involved in peroxisomal proliferation and may regulate peroxisomal membrane., PTM:Seems not to be N-glycosylated, similarity: Belongs to the peroxisomal amembrane., PTM:Seems not to be N-glycosylated, similarity: Belongs to the peroxisomal amembrane and proliferation of peroxisome maintenance and proliferation factors involved in regulation of peroxisome maintenance and proliferation in factors involved in regulation of peroxisome maintenance and proliferation. This gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisoma and membrane and proliferation. This gene protein in 9 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Specificity	PX11A Polyclonal Antibody detects endogenous levels of protein.
ControlThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCollegeObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisome division. May mediate binding of coatomer proteins to the peroxisome at invity. Subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome addition factors involved in corease peroxisome and membrane elongation factors involved in peroxisomal membrane protein to the peroxisomeal division. May mediate binding of coatomer proteins to the peroxisome addition factors involved in peroxisomal proliferation and may regulate peroxisomal membrane. PTM: Seems not to be N-glycosylated, similarity: Belongs to the peroxisonal membrane of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome addition and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisome division. May mediate binding of coatomer proteins to the peroxisomal membrane.PTM: Seems not to be N-glycosylated., similarity: Belongs similarity). Interacts with PEX19., Submit: Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19., and may regulate elongation factors involved in regulation of peroxisome and membrane elongation factors involved in regulation of peroxisome and membrane elongation factors involved in regulation of peroxisome and membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 27kD Cell Pathway Peroxisome membrane ; Multi-pass membrane protein . Tissue Specificity Breast,Liver,Lung,PCR rescued clones, Function function:May be involved in peroxisomal proliferation and may regulate peroxisomed membrane.,PTM:Seems not to be N-glycosylated,similarity:Belongs to the peroxin-11 family, subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19., Background This gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 27kD Cell Pathway Peroxisome membrane ; Multi-pass membrane protein . Tissue Specificity Breast,Liver,Lung,PCR rescued clones, Function function:May be involved in peroxisomal proliferation and may regulate peroxisomed division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxisomal membrane of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Concentration	1 mg/ml
SynonymsObserved Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane ,PTM:Seems not to be N-glycosylated, similarity:Belongs to the peroxin-11 family, subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Purity	≥90%
Observed Band27kDCell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19,,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Storage Stability	-20°C/1 year
Cell PathwayPeroxisome membrane ; Multi-pass membrane protein .Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Synonyms	
Tissue SpecificityBreast,Liver,Lung,PCR rescued clones,Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Observed Band	27kD
Functionfunction:May be involved in peroxisomal proliferation and may regulate peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Cell Pathway	Peroxisome membrane ; Multi-pass membrane protein .
peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By similarity). Interacts with PEX19.,BackgroundThis gene is a member of the PEX11 family, which is composed of membrane elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Tissue Specificity	Breast,Liver,Lung,PCR rescued clones,
elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance. Alternatively spliced transcript variants encoding multiple isoforms have been	Function	peroxisomes division. May mediate binding of coatomer proteins to the peroxisomal membrane.,PTM:Seems not to be N-glycosylated.,similarity:Belongs to the peroxin-11 family.,subunit:Probably interacts with COPB2 and COPA (By
Naniing BYabscience technology CoLtd	Background	elongation factors involved in regulation of peroxisome maintenance and proliferation. This gene product interacts with peroxisomal membrane protein 19 and may respond to outside stimuli to increase peroxisome abundance.
······································		Nanjing BYabscience technology Co.,Ltd






observed for this gene. [provided by RefSeq, Oct 2012],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd