



## TTF2 Polyclonal Antibody

Catalog No	BYab-06307
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	TTF2
Protein Name	Transcription termination factor 2 (EC 3.6.4) (Lodestar homolog) (RNA polymerase II termination factor) (Transcription release factor 2) (F2) (HuF2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TTF2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	127kD
Cell Pathway	Cytoplasm . Nucleus . Cytoplasmic during interphase. Relocates to the nucleus as cells enter mitosis.
Tissue Specificity	Aorta endothelial cell,Brain,Epithelium,
Function	function:DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. May contribute to mitotic transcription repression. May also be involved in pre-mRNA splicing.,similarity:Belongs to the SNF2/RAD54 helicase family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,subcellular location:Cytoplasmic during interphase. Relocates to the nucleus as cells enter mitosis.,subunit:Interacts with CDC5L. Part of the spliceosome.,
Background	This gene encodes a member of the SWI2/SNF2 family of proteins, which play a critical role in altering protein-DNA interactions. The encoded protein has been shown to have dsDNA-dependent ATPase activity and RNA polymerase II
	Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658



国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询

	termination activity. This protein interacts with cell division cycle 5-like, associates with human splicing complexes, and plays a role in pre-mRNA splicing. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images** 

Nanjing BYabscience technology Co.,Ltd