



## RNF10 rabbit pAb

Catalog No	BYab-11807
Isotype	lgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	RNF10 KIAA0262 RIE2
Protein Name	RNF10
Immunogen	Synthesized peptide derived from human RNF10 AA range: 294-344
Specificity	This antibody detects endogenous levels of RNF10 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year Cytoplasm . Nucleus .
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	
Synonyms Observed Band Cell Pathway Tissue Specificity	Cytoplasm . Nucleus . similarity:Belongs to the RNF10 family.,similarity:Contains 1 RING-type zinc
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Cytoplasm . Nucleus . similarity:Belongs to the RNF10 family.,similarity:Contains 1 RING-type zinc finger.,subunit:Interacts with MEOX2., The protein encoded by this gene contains a ring finger motif, which is known to be involved in protein-protein interactions. The specific function of this protein has not yet been determined. EST data suggests the existence of multiple alternatively spliced transcript variants, however, their full length nature is not

## Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co., Ltd