



PTCD3 rabbit pAb

Catalog No	BYab-11850
lsotype	lgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	PTCD3 TRG15
Protein Name	PTCD3
Immunogen	Synthesized peptide derived from human PTCD3 AA range: 624-674
Specificity	This antibody detects endogenous levels of PTCD3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
0	
Concentration	1 mg/ml
Concentration Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Mitochondrion .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Mitochondrion . Abundant in testes, skeletal muscle and heart tissue. similarity:Belongs to the PTCD3 family.,similarity:Contains 10 PPR
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Mitochondrion . Abundant in testes, skeletal muscle and heart tissue. similarity:Belongs to the PTCD3 family.,similarity:Contains 10 PPR
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Mitochondrion . Abundant in testes, skeletal muscle and heart tissue. similarity:Belongs to the PTCD3 family.,similarity:Contains 10 PPR (pentatricopeptide) repeats.,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd