



# FBLI1 rabbit pAb

<b>Catalog No</b>	BYab-12055
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	FBLIM1 FBLP1
<b>Protein Name</b>	FBLI1
<b>Immunogen</b>	Synthesized peptide derived from human FBLI1 AA range: 157-207
<b>Specificity</b>	This antibody detects endogenous levels of FBLI1 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cell junction, focal adhesion . Cytoplasm, cytoskeleton, stress fiber . Associated with actin stress fiber at cell-ECM focal adhesion sites (PubMed:12679033, PubMed:18829455). Isoform 1 and isoform 3 are recruited and localized at actin stress fibers and clustered at cell-EMC adhesion sites through interaction with FERMT2 (PubMed:12679033). Isoform 2 is localized at actin stress fibers (PubMed:12496242). .
<b>Tissue Specificity</b>	Isoform 1 and isoform 3 are expressed in heart, kidney, lung, pancreas, placenta and platelets. Isoform 2 is expressed in brain, heart, kidney, lung, pancreas, placenta, skeletal muscle and platelets.
<b>Function</b>	function:Serves as an anchoring site for cell-ECM adhesion proteins and filamin-containing actin filaments. Is implicated in cell shape modulation (spreading) and motility. May participate in the regulation of filamin-mediated cross-linking and stabilization of actin filaments. May also regulate the assembly of filamin-containing signaling complexes that control actin assembly.,similarity:Contains 3 LIM zinc-binding domains.,subcellular

**Nanjing BYabscience technology Co.,Ltd**



location:Associated with actin stress fiber at cell-ECM focal adhesion sites. Isoform 1 and isoform 3 are recruited and localized at actin stress fibers and clustered at cell-ECM adhesion sites through interaction with PLEKHC1. Isoform 2 is localized at actin stress fibers.,subunit:Isoform 1 and isoform 3 interact with PLEKHC1, FLNA and FLNC. Isoform 2 interacts with FLNB. Interacts with NKX2-5.,tissue specificity:Isoform 1 and isoform 3 are expressed in

#### Background

This gene encodes a protein with an N-terminal filamin-binding domain, a central proline-rich domain, and, multiple C-terminal LIM domains. This protein localizes at cell junctions and may link cell adhesion structures to the actin cytoskeleton. This protein may be involved in the assembly and stabilization of actin-filaments and likely plays a role in modulating cell adhesion, cell morphology and cell motility. This protein also localizes to the nucleus and may affect cardiomyocyte differentiation after binding with the CSX/NKX2-5 transcription factor. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],


#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658