



## RABEK rabbit pAb

Catalog No	BYab-12331
Isotype	lgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	RABEPK RAB9P40
Protein Name	RABEK
Immunogen	Synthesized peptide derived from human RABEK AA range: 227-277
Specificity	This antibody detects endogenous levels of RABEK at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm. Endosome membrane . Interaction with PIKFYVE and subsequent phosphorylation recruits it to the endosomal membrane
Tissue Specificity	
Function	function:Rab9 effector required for endosome to trans-Golgi network (TGN) transport.,PTM:Phosphorylated on Ser residues most probably by PIP5K3.,similarity:Contains 5 Kelch repeats.,subcellular location:Interaction with PIP5K3 and subsequent phosphorylation recruits it to the endosomal membrane.,subunit:Interacts with PIP5K3. Interacts with RAB9 in its GTP-bound conformation.,
Background	
matters needing attention	Avoid repeated freezing and thawing!
	Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




## **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd