



Syntenin mouse mAb

Catalog No	BYab-12625
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	sdcbp
Protein Name	
Immunogen	Purified recombinant human Syntenin protein fragments expressed in E.coli.
Specificity	This antibody detects endogenous levels of Syntenin and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:1000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MDA-9;MDA9;Melanoma differentiation-associated protein 9;Pro-TGF-alpha cytoplasmic domain-interacting protein 18;Scaffold protein Pbp1;SDCB1_HUMAN;Sdcbp;SDCBP;ST1;SYCL; Syndecan binding protein (syntenin);Syndecan binding protein 1;Syndecan-binding protein 1;Syntenin 1;Syntenin-1;TACIP18.
Observed Band	32kD
Cell Pathway	Cell junction, focal adhesion . Cell junction, adherens junction . Cell membrane ; Peripheral membrane protein . Endoplasmic reticulum membrane ; Peripheral membrane protein . Nucleus . Melanosome . Cytoplasm, cytosol . Cytoplasm, cytoskeleton . Secreted, extracellular exosome . Membrane raft . Mainly membrane-associated. Localized to adherens junctions, focal adhesions and endoplasmic reticulum. Colocalized with actin stress fibers. Also found in the nucleus. Identified by mass spectrometry in melanosome fractions from stage I to stage IV. Associated to the plasma membrane in the presence of FZD7 and phosphatidylinositol 4,5-bisphosphate (PIP2) (PubMed:27386966). .
Tissue Specificity	Expressed in lung cancers, including adenocarcinoma, squamous cell carcinoma and small-cell carcinoma (at protein level) (PubMed:25893292). Widely

Nanjing BYabscience technology Co.,Ltd



expressed. Expressed in fetal kidney, liver, lung and brain. In adult highest expression in heart and placenta.

Function	<p>function: Seems to function as an adapter protein. In adherens junctions may function to couple syndecans to cytoskeletal proteins or signaling components. Seems to couple transcription factor SOX4 to the IL-5 receptor (IL5RA). May also play a role in vesicular trafficking. Seems to be required for the targeting of TGFA to the cell surface in the early secretory pathway., induction: By gamma interferon in melanoma cells., PTM: Phosphorylated on tyrosine residues., similarity: Contains 2 PDZ (DHR) domains., subcellular location: Mainly membrane-associated. Localized to adherens junctions, focal adhesions and endoplasmic reticulum. Colocalized with actin stress fibers. Also found in the nucleus. Identified by mass spectrometry in melanosome fractions from stage I to stage IV., subunit: Monomer and homodimer (By similarity). Interacts with SDC1, SDC2, SDC3, SDC4, NRXN2, EPHA7, EPHB1, NF2 isoform 1, TG</p>
Background	<p>syndecan binding protein(SDCBP) Homo sapiens The protein encoded by this gene was initially identified as a molecule linking syndecan-mediated signaling to the cytoskeleton. The syntenin protein contains tandemly repeated PDZ domains that bind the cytoplasmic, C-terminal domains of a variety of transmembrane proteins. This protein may also affect cytoskeletal-membrane organization, cell adhesion, protein trafficking, and the activation of transcription factors. The protein is primarily localized to membrane-associated adherens junctions and focal adhesions but is also found at the endoplasmic reticulum and nucleus. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008],</p>
matters needing attention	<p>Avoid repeated freezing and thawing!</p>
Usage suggestions	<p>This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.</p>

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658



Products Images



Western blot detection of Syntenin in HeLaNE and HeLa cell lysates using Syntenin mouse mAb (1:1000 diluted). Predicted band size: 32KDa. Observed band size: 32KDa.