



ACE1 Polyclonal Antibody

Catalog No	BYab-02466
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	ACE
Protein Name	Angiotensin-converting enzyme
Immunogen	The antiserum was produced against synthesized peptide derived from human ACE1. AA range:891-940
Specificity	ACE1 Polyclonal Antibody detects endogenous levels of ACE1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ACE; DCP; DCP1; Angiotensin-converting enzyme; ACE; Dipeptidyl carboxypeptidase I; Kininase II; CD antigen CD143
Observed Band	165kD
Cell Pathway	[Angiotensin-converting enzyme, soluble form]: Secreted.; Cell membrane; Single-pass type I membrane protein. Cytoplasm . Detected in both cell membrane and cytoplasm in neurons. .
Tissue Specificity	Ubiquitously expressed, with highest levels in lung, kidney, heart, gastrointestinal system and prostate. Isoform Testis-specific is expressed in spermatocytes and adult testis.
Function	catalytic activity:Release of a C-terminal dipeptide, oligopeptide-[Xaa-Yaa, when Xaa is not Pro, and Yaa is neither Asp nor Glu. Thus, conversion of angiotensin I to angiotensin II, with increase in vasoconstrictor activity, but no action on angiotensin II.;cofactor:Bounds 2 zinc ions per subunit. The Testis-specific isoform only binds 1 zinc ion per subunit.;cofactor:Bounds 3 chloride ions per subunit.;disease:Defects in ACE are a cause of renal tubular dysgenesis (RTD) [MIM:267430]. RTD is an autosomal recessive severe disorder of renal tubular development characterized by persistent fetal anuria and perinatal death,

Nanjing BYabscience technology Co.,Ltd



probably due to pulmonary hypoplasia from early-onset oligohydramnios (the Potter phenotype).,disease:Genetic variations in ACE could influence susceptibility to diabetic nephropathy [MIM:612624]; also called susceptibility to microvascular complications of diabetes type 3

Background

This gene encodes an enzyme involved in catalyzing the conversion of angiotensin I into a physiologically active peptide angiotensin II. Angiotensin II is a potent vasopressor and aldosterone-stimulating peptide that controls blood pressure and fluid-electrolyte balance. This enzyme plays a key role in the renin-angiotensin system. Many studies have associated the presence or absence of a 287 bp Alu repeat element in this gene with the levels of circulating enzyme or cardiovascular pathophysiologies. Multiple alternatively spliced transcript variants encoding different isoforms have been identified, and two most abundant spliced variants encode the somatic form and the testicular form, respectively, that are equally active. [provided by RefSeq, May 2010],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




Products Images



Western Blot analysis of HuvEc cells using ACE1 Polyclonal Antibody diluted at 1:1000



Immunohistochemistry analysis of ACE1 antibody in paraffin-embedded human brain tissue.



Western blot analysis of lysates from mouse kidney, using ACE1 antibody.