



# DUS2L Polyclonal Antibody

<b>Catalog No</b>	BYab-02625
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	DUS2L
<b>Protein Name</b>	tRNA-dihydrouridine(20) synthase [NAD(P)+]-like
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human DUS2L. AA range:421-470
<b>Specificity</b>	DUS2L Polyclonal Antibody detects endogenous levels of DUS2L protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	DUS2L; DUS2; tRNA-dihydrouridine(20) synthase [NAD(P)+]-like; Up-regulated in lung cancer protein 8; URLC8; tRNA-dihydrouridine synthase 2-like; hDUS2
<b>Observed Band</b>	55kD
<b>Cell Pathway</b>	Cytoplasm . Endoplasmic reticulum . Mainly at the endoplasmic reticulum. .
<b>Tissue Specificity</b>	Weak expression in heart, placenta and skeletal muscle. Up-regulated in most lung cancer cells (at protein level).
<b>Function</b>	cofactor:FAD.,function:Dihydrouridine synthase. Catalyzes the synthesis of dihydrouridine, a modified base found in the D-loop of most tRNAs.,similarity:Belongs to the dus family. Dus2 subfamily.,similarity:Contains 1 DRBM (double-stranded RNA-binding) domain.,subcellular location:Mainly at the endoplasmic reticulum.,subunit:Interacts with EPRS.,tissue specificity:Weak expression in heart, placenta and skeletal muscle. Up-regulated in most lung cancer cells (at protein level).,
<b>Background</b>	dihydrouridine synthase 2(DUS2) Homo sapiens This gene encodes a cytoplasmic protein that catalyzes the conversion of uridine residues to

**Nanjing BYabscience technology Co.,Ltd**



dihydrouridine in the D-loop of tRNA. The resulting modified bases confer enhanced regional flexibility to tRNA. The encoded protein may increase the rate of translation by inhibiting an interferon-induced protein kinase. This gene has been implicated in pulmonary carcinogenesis. Alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Nov 2012],


**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemistry analysis of paraffin-embedded human skeletal muscle tissue, using DUS2L Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from RAW264.7 cells, using DUS2L Antibody. The lane on the right is blocked with the synthesized peptide.