



## Gab 1 (phospho Tyr627) Polyclonal Antibody

| Catalog No         | BYab-03518  |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human;Mouse;Monkey  |
| Applications       | WB;ELISA  |
| Gene Name          | GAB1  |
| Protein Name       | GRB2-associated-binding protein 1   |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human GAB1 around the phosphorylation site of Tyr627 AA range:611-643   |
| Specificity        | Phospho-Gab 1 (Y627) Polyclonal Antibody detects endogenous levels of Gab 1 protein only when phosphorylated at Y627.   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Polyclonal, Rabbit,IgG  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | GAB1; GRB2-associated-binding protein 1; GRB2-associated binder 1; Growth factor receptor bound protein 2-associated protein 1  |
| Observed Band      | 110kD   |
| Cell Pathway       | cytosol,  |
| Tissue Specificity | Testis,Whole embryo,  |
| Function           | function:Probably involved in EGF and insulin receptor signaling.,PTM:Phosphorylated on tyrosine residue(s) by the epidermal growth factor receptor (EGFR) and the insulin receptor (INSR). Tyrosine phosphorylation of GAB1 mediates interaction with several proteins that contain SH2 domains.,similarity:Belongs to the GAB family.,similarity:Contains 1 PH domain.,subunit:Interacts with GRB2 and with other SH2-containing proteins. Interacts with phosphorylated LAT2., |
| Background         | GRB2 associated binding protein 1(GAB1) Homo sapiens The protein encoded by this gene is a member of the IRS1-like multisubstrate docking protein   |
|                    |   |

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






|                           | family. It is an important mediator of branching tubulogenesis and plays a central role in cellular growth response, transformation and apoptosis. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing!  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.   |

## **Products Images**



Western Blot analysis of various cells using Phospho-Gab 1 (Y627) Polyclonal Antibody



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using GAB1 (Phospho-Tyr627) Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



