



## TCTP (phospho-Ser46) rabbit pAb

Catalog No         BYab-03646           Isotype         IgG           Reactivity         Human;Mouse;Rat           Applications         WB           Gene Name         TPT1           Protein Name         TCTP (Ser46)           Immunogen         Synthesized phosho peptide around human TCTP (Ser46)           Specificity         This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.           Dilution         WB 1:1000-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)           Observed Band         24kD           Cell Pathway         Cytoplasm .           Tissue Specificity         Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and p		
Reactivity  Applications  WB  Gene Name  TPT1  Protein Name  TCTP (Ser46)  Immunogen  Synthesized phosho peptide around human TCTP (Ser46)  Specificity  This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)  Formulation  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution  WB 1:1000-2000  Concentration  1 mg/ml  Purity  290%  Storage Stability  -20°C/1 year  Synonyms  Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band  24kD  Cell Pathway  Cytoplasm  Tissue Specificity  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and Jumphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  WB 1:200-2000  Function: Involved in calcium binding and microtubule stabilization , similarity. Belongs to the TCTP family, subunit: Interacts with STEAP3, tissue specificity-Found in several healthy and tumoral cells including erythrocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and Jumphomass. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.	Catalog No	BYab-03646
Applications WB Gene Name TPT1 Protein Name TCTP (Ser46) Immunogen Synthesized phosho peptide around human TCTP (Ser46) Specificity This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46) Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dillution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23) Observed Band 24kD Cell Pathway Cytoplasm . Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate. Function light and renal cell carcinoma (RCC). Expressed in lealthy and tumoral cells including erythrocytes, hepatocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.	Isotype	IgG
Gene Name TPT1 Protein Name TCTP (Ser46)  Immunogen Synthesized phosho peptide around human TCTP (Ser46)  Specificity This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1:1000-2000  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band 24kD  Cell Pathway Cytoplasm .  Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function Individual calcium binding and microubbule stabilization, similarity. Belongs to the TCTP family, subunit: Interacts with STEAP3, tissue specificity. Found in several healthy and tumoral cells including erythrocytes, hepatocytes, nepatocytes, ne	Reactivity	Human;Mouse;Rat
Protein Name TCTP (Ser46)  Immunogen Synthesized phosho peptide around human TCTP (Ser46)  Specificity This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1:1000-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band 24kD  Cell Pathway Cytoplasm .  Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3, tissue specificity:Found in several healthy and tumoral cells including erythrocytes, negrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Applications	WB
Immunogen         Synthesized phosho peptide around human TCTP (Ser46)           Specificity         This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.           Dilution         WB 1:1000-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)           Observed Band         24kD           Cell Pathway         Cytoplasm           Tissue Specificity         Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.           Function         function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAPS, itssue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be d	Gene Name	TPT1
Specificity         This antibody detects endogenous levels of (phospho-Ser46)           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.           Dilution         WB 1:1000-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)           Observed Band         24kD           Cell Pathway         Cytoplasm .           Tissue Specificity         Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.           Function         function: Involved in calcium binding and microtubule stabilization , similarity. Belongs to the TCTP family , subunit: Interacts with STEAP3, tissue specificity: Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Protein Name	TCTP (Ser46)
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1:1000-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band 24kD  Cell Pathway Cytoplasm .  Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family, subunit:Interacts with STEAP3. tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.	Immunogen	Synthesized phosho peptide around human TCTP (Ser46)
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       WB 1:1000-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)         Observed Band       24kD         Cell Pathway       Cytoplasm .         Tissue Specificity       Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.         Function       function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family, subunit:Interacts with STEAP3, tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Specificity	This antibody detects endogenous levels of Human Mouse Rat TCTP (phospho-Ser46)
Purification  The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution  WB 1:1000-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band  24kD  Cell Pathway  Cytoplasm  Tissue Specificity  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinomy and microtubule stabilization., similarity:Belongs to the TCTP family., subunit:Interacts with STEAP3., tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen.  Dilution WB 1:1000-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band 24kD  Cell Pathway Cytoplasm .  Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)         Observed Band       24kD         Cell Pathway       Cytoplasm .         Tissue Specificity       Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.         Function       function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family,,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band 24kD  Cell Pathway Cytoplasm .  Tissue Specificity Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Dilution	WB 1:1000-2000
Storage Stability  -20°C/1 year  Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band  24kD  Cell Pathway  Cytoplasm.  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Concentration	1 mg/ml
Synonyms  Translationally-controlled tumor protein (TCTP) (Fortilin) (Histamine-releasing factor) (HRF) (p23)  Observed Band  24kD  Cell Pathway  Cytoplasm .  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Purity	≥90%
Cell Pathway  Cytoplasm .  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm .  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Synonyms	
Tissue Specificity  Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Observed Band	24kD
macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in placenta and prostate.  Function  function:Involved in calcium binding and microtubule stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Cell Pathway	Cytoplasm .
stabilization.,similarity:Belongs to the TCTP family.,subunit:Interacts with STEAP3.,tissue specificity:Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in	Tissue Specificity	macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney
	Function	stabilization., similarity: Belongs to the TCTP family., subunit: Interacts with STEAP3., tissue specificity: Found in several healthy and tumoral cells including erythrocytes, hepatocytes, macrophages, platelets, keratinocytes, erythroleukemia cells, gliomas, melanomas, hepatoblastomas, and lymphomas. It cannot be detected in kidney and renal cell carcinoma (RCC). Expressed in

Nanjing BYabscience technology Co.,Ltd






matters	needing	
attention		

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658