



IGKC Polyclonal Antibody

Catalog No BYab-07747 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name IGKC Protein Name Ig kappa chain C region Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Leu-83, similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background Mydi-B3. The ROY se		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name IGKC Protein Name Ig kappa chain C region Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous: The EU sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83. similarity: Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-07747
Applications WB;ELISA Gene Name IGKC Protein Name Ig kappa chain C region Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name IGKC Protein Name Ig kappa chain C region Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic	Reactivity	Human;Rat;Mouse;
Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity:Contains 1 Ig-like (immunoglobulin-like) domain., miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (3) allotypic marker, Ala-45 and Leu-83., similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Gene Name	IGKC
Specificity IGKC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous: The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity: Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Protein Name	Ig kappa chain C region
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous: The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity: Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Matters needing Avoid repeated freezing and thawing!	Specificity	IGKC Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. Similarity:Contains 1 Ig-like (immunoglobulin-like) domain.	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 lg-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 lg-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous: The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity: Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous: The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83., similarity: Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band 11kD Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Secreted . Cell membrane . Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Abdominal adipose tissue, Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Observed Band	11kD
Function miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Cell Pathway	Secreted . Cell membrane .
Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., Background miscellaneous:The EU sequence has the INV (3) allotypic marker, Ala-45 and Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Abdominal adipose tissue,
Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and Leu-83.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain., matters needing Avoid repeated freezing and thawing!	Function	Val-83. The ROY sequence has the INV (1,2) allotypic marker, Ala-45 and
	Background	Val-83. The ROY sequence has the INV (1.2) allotypic marker. Ala-45 and
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

