





MIEN1 Polyclonal Antibody

Catalog No	BYab-07848
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	MIEN1 C17orf37 RDX12 XTP4
Protein Name	Migration and invasion enhancer 1 (HBV X-transactivated gene 4 protein) (HBV XAg-transactivated protein 4) (Protein C35)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Specificity	MIEN1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year 12kD Cytoplasm, cytosol. Cell membrane; Lipid-anchor; Cytoplasmic side.
Storage Stability Synonyms Observed Band	-20°C/1 year 12kD Cytoplasm, cytosol. Cell membrane; Lipid-anchor; Cytoplasmic side. Concentrates at the leading edge of migrating cells. Localizes outside membrane
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 12kD Cytoplasm, cytosol. Cell membrane; Lipid-anchor; Cytoplasmic side. Concentrates at the leading edge of migrating cells. Localizes outside membrane raft regions. Among normal tissues, present only in Leydig cells. Strongly up-regulated in breast cancers and in brain cancer distant metastasis (at protein level). Up-regulated in prostate cancer cells and in the higher grades of prostate
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 12kD Cytoplasm, cytosol. Cell membrane; Lipid-anchor; Cytoplasmic side. Concentrates at the leading edge of migrating cells. Localizes outside membrane raft regions. Among normal tissues, present only in Leydig cells. Strongly up-regulated in breast cancers and in brain cancer distant metastasis (at protein level). Up-regulated in prostate cancer cells and in the higher grades of prostate adenocarcinoma (at protein level). function:May be involved in a redox-related process .,similarity:Belongs to the
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year 12kD Cytoplasm, cytosol. Cell membrane; Lipid-anchor; Cytoplasmic side. Concentrates at the leading edge of migrating cells. Localizes outside membrane raft regions. Among normal tissues, present only in Leydig cells. Strongly up-regulated in breast cancers and in brain cancer distant metastasis (at protein level). Up-regulated in prostate cancer cells and in the higher grades of prostate adenocarcinoma (at protein level). function:May be involved in a redox-related process .,similarity:Belongs to the SelWTH family., function:May be involved in a redox-related process .,similarity:Belongs to the

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

