



## 5NT3L rabbit pAb

Catalog No     BYab-08456       Isotype     IgG       Reactivity     Human; Mouse;Rat       Applications     WB       Gene Name     NT5C3L       Protein Name     5NT3L       Immunogen     Synthesized peptide derived from human 5NT3L AA range: 23-73       Specificity     This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat       Formulation     Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.       Source     Polyclonal, Rabbit,IgG       Purification     The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.       Dilution     WB 1: 500-2000       Concentration     1 mg/ml       Purity     ≥90%       Storage Stability     -20°C/1 year       Synonyms     Observed Band       Cell Pathway     Cytoplasm .       Tissue Specificity       Function     catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate, function: Can act both as nucleotidase and as phosphotransferase, similarity: Belongs to the pyrimidine 5'-nucleotidase familly.       Background     Avoid repeated freezing and thawing!		
Reactivity Human; Mouse;Rat  Applications WB  Gene Name NT5C3L  Protein Name 5NT3L  Immunogen Synthesized peptide derived from human 5NT3L AA range: 23-73  Specificity This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1: 500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasm .  Tissue Specificity  Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate, function:Can act both as nucleotidase and as phosphotransferase, similarity: Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-08456
Applications  WB  Gene Name NT5C3L  Protein Name SNT3L  Immunogen Synthesized peptide derived from human 5NT3L AA range: 23-73  Specificity This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1: 500-2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band Cell Pathway Cytoplasm.  Tissue Specificity  Function catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name NT5C3L Protein Name 5NT3L Immunogen Synthesized peptide derived from human 5NT3L AA range: 23-73 Specificity This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band Cell Pathway Cytoplasm . Tissue Specificity Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate, function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family., Background  matters needing Avoid repeated freezing and thawing!	Reactivity	Human; Mouse;Rat
Protein Name 5NT3L  Immunogen Synthesized peptide derived from human 5NT3L AA range: 23-73  Specificity This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1: 500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasm .  Tissue Specificity  Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate, function: Can act both as nucleotidase and as phosphotransferase, similarity: Belongs to the pyrimidine 5'-nucleotidase family.  Background  matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen       Synthesized peptide derived from human 5NT3L AA range: 23-73         Specificity       This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       WB 1: 500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasm .         Tissue Specificity         Function       catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate, function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family.,         Background       Avoid repeated freezing and thawing!	Gene Name	NT5C3L
Specificity       This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       WB 1: 500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasm .         Tissue Specificity       Function       catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,         Background       Avoid repeated freezing and thawing!	Protein Name	5NT3L
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution WB 1: 500-2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band Cell Pathway Cytoplasm.  Tissue Specificity Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate., function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family.,  Background matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human 5NT3L AA range: 23-73
Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       WB 1: 500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasm .         Tissue Specificity       Function       catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate_function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family.,         Background       Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of 5NT3L at Human/Mouse/Rat
Purification  The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution  WB 1: 500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  Cell Pathway  Cytoplasm.  Tissue Specificity  Function  catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function: Can act both as nucleotidase and as phosphotransferase.,similarity: Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen.  Dilution WB 1: 500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasm.  Tissue Specificity  Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasm .  Tissue Specificity  Function catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasm .  Tissue Specificity  Function catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate., function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band  Cell Pathway Cytoplasm .  Tissue Specificity  Function catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasm .  Tissue Specificity  Function  catalytic activity: A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate., function: Can act both as nucleotidase and as phosphotransferase., similarity: Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasm .  Tissue Specificity  Function  catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing  Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity  Function  catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  matters needing  Avoid repeated freezing and thawing!	Observed Band	
Function  catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  Avoid repeated freezing and thawing!	Cell Pathway	Cytoplasm .
phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,  Background  Matters needing  Avoid repeated freezing and thawing!	Tissue Specificity	
matters needing Avoid repeated freezing and thawing!	Function	catalytic activity:A 5'-ribonucleotide + H(2)O = a ribonucleoside + phosphate.,function:Can act both as nucleotidase and as phosphotransferase.,similarity:Belongs to the pyrimidine 5'-nucleotidase family.,
matters needing Avoid repeated freezing and thawing!	Background	
	matters needing attention	Avoid repeated freezing and thawing!

## Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd