



CJ099 rabbit pAb

Catalog No	BYab-08657
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF
Gene Name	C10orf99 UNQ1833/PRO3446
Protein Name	CJ099
Immunogen	Synthesized peptide derived from human CJ099 AA range: 19-69
Specificity	This antibody detects endogenous levels of CJ099 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Secreted . Expressed at high levels in colon, and cervix and at moderate level in tonsil (PubMed:25585381, PubMed:28900043). Highly reduced expression in primary colon cancer tissues compared with that in adjacent tissues (PubMed:25351403). Highest levels of expression detected in stomach and colon; expressed in epithelium of skin and esophagus, and in some tumor and/or tumor adjacent tissues (TAT), including TAT of esophagus cancer, hepatocellular carcinoma
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Secreted . Expressed at high levels in colon, and cervix and at moderate level in tonsil (PubMed:25585381, PubMed:28900043). Highly reduced expression in primary colon cancer tissues compared with that in adjacent tissues (PubMed:25351403). Highest levels of expression detected in stomach and colon; expressed in epithelium of skin and esophagus, and in some tumor and/or tumor adjacent tissues (TAT), including TAT of esophagus cancer, hepatocellular carcinoma (HCC), squamous cell carcinoma (SCC), basal cell carcinoma (BCC) and invasive

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Nanjing BYabscience technology Co.,Ltd