



MEST rabbit pAb

| Catalog No | BYab-09452 |
|--|--|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | MEST PEG1 |
| Protein Name | Mesoderm-specific transcript homolog protein (EC 3) (Paternally-expressed gene 1 protein) |
| Immunogen | Synthesized peptide derived from human MEST AA range: 187-237 |
| Specificity | This antibody detects endogenous levels of human MEST |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:1000-2000 |
| | |
| Concentration | 1 mg/ml |
| Concentration Purity | 1 mg/ml ≥90% |
| | • |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year Endoplasmic reticulum membrane; Multi-pass membrane protein. Highly expressed in hydatidiform moles, but barely expressed in dermoid cysts. Biallelic expression is detected in blood lymphocytes. Seems to imprinted in an isoform-specific manner rather than in a tissue-specific manner in lymphocytes. Isoform 1 is expressed only from the paternal allele. Isoform 2 is expressed from |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year Endoplasmic reticulum membrane; Multi-pass membrane protein. Highly expressed in hydatidiform moles, but barely expressed in dermoid cysts. Biallelic expression is detected in blood lymphocytes. Seems to imprinted in an isoform-specific manner rather than in a tissue-specific manner in lymphocytes. Isoform 1 is expressed only from the paternal allele. Isoform 2 is expressed from |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | ≥90% -20°C/1 year Endoplasmic reticulum membrane ; Multi-pass membrane protein . Highly expressed in hydatidiform moles, but barely expressed in dermoid cysts. Biallelic expression is detected in blood lymphocytes. Seems to imprinted in an isoform-specific manner rather than in a tissue-specific manner in lymphocytes. Isoform 1 is expressed only from the paternal allele. Isoform 2 is expressed from |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| ı | Isaq | P S | ua | ae | stic | าทร |
|---|------|------------|----|-----|------|-----|
| u | Jau | C 3 | uu | uc. | วแง | ノロコ |

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

| Products Images | | | | | | |
|-----------------|--|--|--|--|--|--|
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |

Nanjing BYabscience technology Co.,Ltd