



## PDK4 Rabbit pAb

Catalog No	BYab-10306
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	IHC;WB
Gene Name	PDK4 PDHK4
Protein Name	PDK4
Immunogen	Synthesized peptide derived from human PDK4 AA range: 34-84
Specificity	This antibody detects endogenous levels of PDK4 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.03% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification  Dilution	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  IHC-p1:50-200 ,WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	[Pyruvate dehydrogenase [lipoamide]] kinase isozyme 4, mitochondrial (EC 2.7.11.2) (Pyruvate dehydrogenase kinase isoform 4)
Observed Band	46kD
Cell Pathway	Mitochondrion matrix.
Tissue Specificity	Ubiquitous; highest levels of expression in heart and skeletal muscle.
Function	
Background	Pyruvate dehydrogenase kinase isoform4 (PDK4) is also named as PDHK4 and belongs to the PDK/BCKDK protein kinase family. It is upregulated by starvation in many tissues of the body during starvation. This causes inactivation of the pyruvate dehydrogenase complex which blocks pyruvate oxidation and conserves lactate and alanine for gluconeogenesis.
matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询




**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of 1) HepG2 Cell Lysate, 2) Mouse Heart Tissue Lysate, 3)Rat Muscle Tissue Lysate using Rabbit pAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Nanjing BYabscience technology Co.,Ltd