



CDX2 (Phospho-Ser283) rabbit pAb

Mouse,Rat Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.194% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelial lining of both the sma and large intestine. PTM:Phosphorylation of Ser-60 mediates the transactiva		
ReactivityHuman; Mouse;RatApplicationsWBGene NameCDX2 CDX3Protein NameCDX2 (Phospho-Ser283)ImmunogenSynthesized peptide derived from human CDX2 (Phospho-Ser283)SpecificityThis antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Hu Mouse,RatFormulationLiquid in PBS containing 50% glycerol, and 0.194% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandClause.Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early and large intestine. PTM:Phosphorylation of Ser-60 mediates the transactiva and and regine to the smacetiva and and regine to thin of Ser-60 mediates the transactiva and and regine to the smacetiva and and regine to the smacetiva and and regine to the smacetiva and and regine to the smacetiva and and regine to the smacetiva and and regine to the smacetiva and and regine to the smacetiva <br< th=""><td>Catalog No</td><td>BYab-10538</td></br<>	Catalog No	BYab-10538
Applications WB Gene Name CDX2 CDX3 Protein Name CDX2 (Phospho-Ser283) Immunogen Synthesized peptide derived from human CDX2 (Phospho-Ser283) Specificity This antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Hu Mouse, Rat Formulation Liquid in PBS containing 50% glycerol, and 0.194% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Nucleus . Tissue Specificity Detected in small intestine, colon and pancreas. Function function-Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early and furge intestine, pothophophorylation of Ser-60 mediates the transactiva and large intestine, PTM.Phophophorylation of Ser-60 mediates the transactiva and large intestine, PTM.Phophophorylation of Ser-60 mediates the transactiva and large intestine and other intestinal epithelial lining of both the sma and large intestine and of the intrestransactivatin	Isotype	lgG
Gene Name CDX2 CDX3 Protein Name CDX2 (Phospho-Ser283) Immunogen Synthesized peptide derived from human CDX2 (Phospho-Ser283) Specificity This antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Hu Mouse, Rat Formulation Liquid in PBS containing 50% glycerol, and 0.194% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early and large intestinePTM:PhosphorVation of Ser-60 mediates the transactiva and large intestinePTM:PhosphorVation of Ser-60 mediates the transactiva	Reactivity	Human; Mouse;Rat
Protein NameCDX2 (Phospho-Ser283)ImmunogenSynthesized peptide derived from human CDX2 (Phospho-Ser283)SpecificityThis antibody detects endogenous levels of Mouse,RatFormulationLiquid in PBS containing 50% glycerol, and 0.194% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early and larce intestine. PTM-Phosphortation of Seried on mediates the transactiva and larce intestine. PTM-Phosphortation of Seried on mediates the transactiva and larce intestine. PTM-Phosphortation of Seried on mediates the transactiva and larce intestine. PTM-Phosphortation of Seried on seried was and seried the intestinal epithelium. Important in broad range of functions from early and larce intestine. PTM-Phosphortation of Seried on mediates the transactiva and larce intestine. PTM-Phosphortation of Seried on detates the transactiva and larce intestine. PTM-Phosphortation of Seried on detates the transactiva and large intestine. PTM-Phosphortation of Seried on detates the transactiva and large intestine. PTM-Phosphortation of Seried on detates the transactiva and large intestine. PTM-Phosphortation of Seried on detates t	Applications	WB
ImmunogenSynthesized peptide derived from human CDX2 (Phospho-Ser283)SpecificityThis antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Hu Mouse,RatFormulationLiquid in PBS containing 50% glycerol, and 0.194% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandCell PathwayTissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed differentiation to maintenance of the intestinal epithelial lining of both the sma and large intestine.PTM.Phosphorylation of Ser-60 mediates the transactiva	Gene Name	CDX2 CDX3
Specificity This antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Hu Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.194% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the sma and larce intestine.PMC Phosphorylation of Seried Berlow	Protein Name	CDX2 (Phospho-Ser283)
Mouse,Rat Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.194% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the smal and large intestine. PTM:Phosphorylation of Ser-60 mediates the transactiva	Immunogen	Synthesized peptide derived from human CDX2 (Phospho-Ser283)
SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandNucleus .Cell PathwayNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactiva	Specificity	This antibody detects endogenous levels of CDX2 (Phospho-Ser283) at Human, Mouse,Rat
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatograph using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelial lining of both the smal and large intestinePTM:Phosphorylation of Ser-60 mediates the transactival and large intestine	Formulation	Liquid in PBS containing 50% glycerol, and 0.194% sodium azide.
using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early addiferentiation to maintenance of the intestinal epithelium ling of both the smal and large intestinePTM:Phosphorylation of Ser-60 mediates the transactival a	Source	Polyclonal, Rabbit,IgG
Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandCell PathwayCell PathwayNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelial lining of both the smal and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Purity ≥90% Storage Stability -20°C/1 year Synonyms Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2) Observed Band Cell Pathway Nucleus . Nucleus . Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Dilution	WB 1:500-2000
Storage Stability-20°C/1 yearSynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandCell PathwayCell PathwayNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Concentration	1 mg/ml
SynonymsHomeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)Observed BandNucleus .Cell PathwayNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Purity	≥90%
Observed Band Cell Pathway Nucleus . Tissue Specificity Detected in small intestine, colon and pancreas. Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Storage Stability	-20°C/1 year
Cell PathwayNucleus .Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Synonyms	Homeobox protein CDX-2 (CDX-3) (Caudal-type homeobox protein 2)
Tissue SpecificityDetected in small intestine, colon and pancreas.Functionfunction:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation	Observed Band	
Function function:Involved in the transciptional regulation of multiple genes expressed the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the sma and large intestinePTM:Phosphorylation of Ser-60 mediates the transactival	Cell Pathway	Nucleus .
the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the sma and large intestinePTM:Phosphorylation of Ser-60 mediates the transactivation ser-60	Tissue Specificity	Detected in small intestine, colon and pancreas.
homeobox DNA-binding domain.,	Function	differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine.,PTM:Phosphorylation of Ser-60 mediates the transactivation capacity.,similarity:Belongs to the Caudal homeobox family.,similarity:Contains 1
family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth an differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gen	Background	This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth an differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis. [provided by RefSeq,
Nanjing BYabscience technology Co.,Ltd		Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Jan 2	2012],
-------	--------

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd