



KCTD6 rabbit pAb

Catalog No	BYab-10967
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	KCTD6
Protein Name	KCTD6
Immunogen	Synthesized peptide derived from human KCTD6 AA range: 118-168
Specificity	This antibody detects endogenous levels of KCTD6 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm, myofibril, sarcomere, M line . Colocalizes with ANK1 isoform Mu17 at the M line in differentiated skeletal muscle cells and heart
Cell Pathway Tissue Specificity	Cytoplasm, myofibril, sarcomere, M line . Colocalizes with ANK1 isoform Mu17 at the M line in differentiated skeletal muscle cells and heart Highly expressed in cerebellum and brain. Expression is down-regulated in medulloblastoma.
	Highly expressed in cerebellum and brain. Expression is down-regulated in
Tissue Specificity	Highly expressed in cerebellum and brain. Expression is down-regulated in medulloblastoma.
Tissue Specificity Function	Highly expressed in cerebellum and brain. Expression is down-regulated in medulloblastoma.
Tissue Specificity Function Background matters needing	medulloblastoma. similarity:Contains 1 BTB (POZ) domain.,


Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






Products Images



Western blot analysis of lysates from A549 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd