



## VEGF-B Polyclonal Antibody

| Catalog No         | BYab-15979  |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human;Mouse;Rat   |
| Applications       | WB;IHC;IF;ELISA   |
| Gene Name          | VEGFB   |
| Protein Name       | Vascular endothelial growth factor B  |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human VEGFB. AA range:91-140  |
| Specificity        | VEGF-B Polyclonal Antibody detects endogenous levels of VEGF-B protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Polyclonal, Rabbit,IgG  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | Western Blot: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000. IF 1:100-300 Not yet tested in other applications.   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           | VEGFB; VRF; Vascular endothelial growth factor B; VEGF-B; VEGF-related factor; VRF  |
| Observed Band      | 21kD  |
| Cell Pathway       | Secreted. Secreted but remains associated to cells or to the extracellular matrix unless released by heparin.   |
| Tissue Specificity | Expressed in all tissues except liver. Highest levels found in heart, skeletal muscle and pancreas.   |
| Function           | alternative products:Additional isoforms seem to exist,function:Growth factor for endothelial cells. VEGF-B167 binds heparin and neuropilin-1 whereas the binding to neuropilin-1 of VEGF-B186 is regulated by proteolysis.,PTM:VEGF-B186 is O-glycosylated.,similarity:Belongs to the PDGF/VEGF growth factor family.,subcellular location:Secreted but remains associated to cells or to the extracellular matrix unless released by heparin.,subunit:Homodimer; disulfide-linked. Can also form heterodimer with VEGF.,tissue specificity:Expressed in all tissues except liver. Highest levels found in heart, skeletal muscle and pancreas., |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




| Background                | This gene encodes a member of the PDGF (platelet-derived growth factor)/VEGF (vascular endothelial growth factor) family. The VEGF family members regulate the formation of blood vessels and are involved in endothelial cell physiology. This member is a ligand for VEGFR-1 (vascular endothelial growth factor receptor 1) and NRP-1 (neuropilin-1). Studies in mice showed that this gene was co-expressed with nuclear-encoded mitochondrial genes and the encoded protein specifically controlled endothelial uptake of fatty acids. Alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Sep 2011], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |







## **Products Images**



Immunofluorescence analysis of A549. 1,primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.




Western Blot analysis of various cells using VEGF-B Polyclonal Antibody diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000




Western Blot analysis of Jurkat cells using VEGF-B Polyclonal Antibody diluted at 1:2000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000


Nanjing BYabscience technology Co.,Ltd









Western blot analysis of lysates from Jurkat cells, using VEGFB Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).







(kD) 117-

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658