



# Gβ 5 Polyclonal Antibody

Catalog No	BYab-16176
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	GNB5
Protein Name	Guanine nucleotide-binding protein subunit beta-5
Immunogen	The antiserum was produced against synthesized peptide derived from human GNB5. AA range:151-200
Specificity	Gβ 5 Polyclonal Antibody detects endogenous levels of Gβ 5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GNB5; Guanine nucleotide-binding protein subunit beta-5; Gbeta5; Transducin beta chain 5
Observed Band	40kD
Cell Pathway	Membrane .
Tissue Specificity	Widely expressed.
Function	function:Guanine nucleotide-binding proteins (G proteins) are involved as a modulator or transducer in various transmembrane signaling systems. The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction.,similarity:Belongs to the WD repeat G protein beta family.,similarity:Contains 7 WD repeats.,subunit:G proteins are composed of 3 units, alpha, beta and gamma. Component of the RGS9-1-Gbeta5 complex composed of RGS9 (isoform RGS9-1), Gbeta5 (GNB5) and RGS9BP.,tissue specificity:Expressed in multiple tissues.,

Nanjing BYabscience technology Co.,Ltd

**Background**

Heterotrimeric guanine nucleotide-binding proteins (G proteins), which integrate signals between receptors and effector proteins, are composed of an alpha, a beta, and a gamma subunit. These subunits are encoded by families of related genes. This gene encodes a beta subunit. Beta subunits are important regulators of alpha subunits, as well as of certain signal transduction receptors and effectors. Alternatively spliced transcript variants encoding different isoforms exist. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of HepG2 cells using Gβ 5 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).