



## SPHK2 Mouse mAb(7A5)

Catalog No	BYab-17215	
Isotype	IgG	
Reactivity	Human, Mouse,Rat	
Applications	WB,IHC	
Gene Name	SPHK2	
Protein Name	SPHK2	
Immunogen	Synthesized peptide derived from human SPHK2	
Specificity	This antibody detects endogenous levels of SPHK2 at Human, Mouse,Rat	
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.	
Source	Mouse,monoclonal	
Purification  Dilution	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 IHC 1:50-200	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms	Sphingosine kinase 2 (SK 2) (SPK 2) (EC 2.7.1.91)	
Observed Band		
Cell Pathway	Cytoplasm . Nucleus . Endoplasmic reticulum . Mitochondrion inner membrane . In nucleus, located in nucleosomes where it associates with core histone proteins such as histone 3 (PubMed:19729656). In brains of patients with Alzheimer's disease, may be preferentially localized in the nucleus. Cytosolic expression decrease correlates with the density of amyloid deposits (PubMed:29615132). In apoptotic cells, colocalizes with CASP1 in cell membrane where is cleaved and released from cells in an active form (PubMed:20197547); [Isoform 2]: Lysosome membrane .	
Tissue Specificity	Mainly expressed in adult kidney, liver, and brain (PubMed:10751414). Expressed in cerebral cortex and hippocampus (at protein level) (PubMed:29615132). Isoform 1 is the predominant form expressed in most tissues (PubMed:16103110).	
Function	alternative products:Experimental confirmation may be lacking for some isoforms,catalytic activity:ATP + sphinganine = ADP + sphinganine 1-phosphate.,catalytic activity:ATP + sphingosine = ADP + sphingosine	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	1-phosphate.,cofactor:Magnesium.,function:Catalyzes the phosphorylation of sphingosine to form sphingosine 1-phosphate (SPP), a lipid mediator with both intra-and extracellular functions. Also acts on D-erythro-dihydrosphingosine, D-erythro-sphingosine and L-threo-dihydrosphingosine.,similarity:Contains 1 DAGKc domain.,	
Background		
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images

Nanjing BYabscience technology Co.,Ltd