



## DDDDK-Tag(binds to flag sequnence) Rabbit mAb(M11)

Catalog No	BYab-17218
Isotype	IgG
Reactivity	Species independent
Applications	WB;ELISA;IF
Gene Name	Flag tag; Flag-tag, DDDDK TAG, DDDDK-TAG, DYKDDDDK tag, DYKDDDK-tag
Protein Name	
Immunogen	Synthetic Peptide of Flag-Tag
Specificity	The antibody detects C-terminal, internal, and N-terminal Flag-tag fusion proteins.
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Rabbit VH IgG1 and VL Kappa was expressed from 293F cells
Purification	Protein A purified
Dilution	WB: 1:1000-5000 ELISA 1:20000-50000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	
Tissue Specificity	
Function	
Background	The DYKDDDK peptide (Flag-tag) is a polypeptide protein tag that can be added to a protein using recombinant DNA technology. It can be used for affinity chromatography, and then used to separate recombinant, over expressed protein from wild-type protein expressed by the host organism. It can also be used in the isolation of protein complexes with multiple subunits.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**




Western Blot analysis of 293F and CHO cells transfected or non-tansfected DDDK-tag expression vector by primary antibody at 1:3000 dilution.



Nanjing BYabscience technology Co.,Ltd







Western Blot analysis of flag-tag protein using primary antibody at various dilution. Secondary antibody(catalog#:RS0002) was diluted at 1:10000

