



Blk (phospho Tyr501) Polyclonal Antibody

Catalog No	BYab-14542
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	BLK
Protein Name	Tyrosine-protein kinase Blk
Immunogen	The antiserum was produced against synthesized peptide derived from human BLK around the phosphorylation site of Tyr501. AA range:456-505
Specificity	Phospho-Blk (Y501) Polyclonal Antibody detects endogenous levels of Blk protein only when phosphorylated at Y501.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/20000 IF 1:50-200
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year BLK; Tyrosine-protein kinase Blk; B lymphocyte kinase; p55-Blk Cell membrane ; Lipid-anchor . Present and active in lipid rafts. Membrane
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year BLK; Tyrosine-protein kinase Blk; B lymphocyte kinase; p55-Blk Cell membrane ; Lipid-anchor . Present and active in lipid rafts. Membrane location is required for the phosphorylation of CD79A and CD79B (By similarity) Expressed in lymphatic organs, pancreatic islets, Leydig cells, striate ducts of
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year BLK; Tyrosine-protein kinase Blk; B lymphocyte kinase; p55-Blk Cell membrane ; Lipid-anchor . Present and active in lipid rafts. Membrane location is required for the phosphorylation of CD79A and CD79B (By similarity) Expressed in lymphatic organs, pancreatic islets, Leydig cells, striate ducts of salivary glands and hair follicles. catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:May function in a signal transduction pathway that is restricted to B-lymphoid cells.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. SRC subfamilysimilarity:Contains 1 protein kinase






[provided by RefSeq, Aug 2010],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using BLK (Phospho-Tyr501) Antibody. The picture on the right is blocked with the phospho peptide.

Nanjing BYabscience technology Co.,Ltd