



# Ki-67 mouse mAb(ABT001)

Catalog No	BYab-15525
Isotype	IgG
Reactivity	Human
Applications	IHC;WB;IF
Gene Name	MKI67
Protein Name	Ki-67
Immunogen	Synthesized peptide derived from human Ki-67
Specificity	The antibody can specifically recognize human Ki-67 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.141% sodium azide.
Source	Mouse, Monoclonal/IgG2b, kappa
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:100-500, WB 1:200-1000, IF 1:100-500
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Antigen KI-67
Observed Band	
Cell Pathway	Chromosome . Nucleus . Nucleus, nucleolus . Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the mitotic chromosome surface (PubMed:27362226). Associates with satellite DNA in G1 phase (PubMed:9510506). Binds tightly to chromatin in interphase, chromatin-binding decreases in mitosis when it associates with the surface of the condensed chromosomes (PubMed:15896774, PubMed:22002106). Predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix (PubMed:22002106). .
Tissue Specificity	Nuclear
Function	developmental stage:Expression of this antigen occurs preferentially during late G1, S, G2 and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected.,function:Thought to be required for maintaining cell proliferation.,online information:Ki-67 entry,similarity:Contains 1 FHA domain.,subcellular location:Predominantly localized in the G1 phase in the

Nanjing BYabscience technology Co.,Ltd



perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix. In mitosis, it is present on all chromosomes.,subunit:Interacts with KIF15. Binds through the FHA domain to MKI67IP.,

**Background**

This gene encodes a nuclear protein that is associated with and may be necessary for cellular proliferation. Alternatively spliced transcript variants have been described. A related pseudogene exists on chromosome X. [provided by RefSeq, Mar 2009],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images




Human appendix tissue was stained with Anti-Ki-67 (ABT001) Antibody



Human colon carcinoma tissue was stained with Anti-Ki-67 (ABT001) Antibody



Human colon carcinoma tissue was stained with Anti-Ki-67 (ABT001) Antibody




Human malignant mesothelioma tissue was stained with Anti-Ki-67 (ABT001) Antibody

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658



Human tonsil tissue was stained with Anti-Ki-67 (ABT001) Antibody